

UT-808

Investigation into $O(N)$ Invariant Scalar Model Using
Auxiliary-Mass Method at Finite Temperature

Kenzo Ogure*

*Institute for Cosmic Ray Research, University of Tokyo, Midori-cho,
Tanashi, Tokyo 188, Japan*

and

Joe Sato†

*Department of Physics, School of Science, University of Tokyo,
Tokyo 113, Japan*

February 1, 2008

Abstract

Using auxiliary-mass method, $O(N)$ invariant scalar model is investigated at finite temperature. This mass and an evolution equation allow us to calculate an effective potential without an infrared divergence. Second order phase transition is indicated by the effective potential. The critical exponents are determined numerically.

*e-mail address: ogure@icrhp3.icrr.u-tokyo.ac.jp

†e-mail address: joe@hep-th.phys.s.u-tokyo.ac.jp

Symmetry restoration of $O(N)$ scalar model at high temperature is very important since many physical systems belong to the same universality class as it: polymer phase transition ($N \rightarrow 0$), critical liquid-vapor phase transition ($N=1$), alloy (e.g. β -brass) phase transition ($N=1$), uniaxial ferromagnet phase transition ($N=1$), superfluid phase transition ($N=2$), ferromagnet phase transition ($N=3$), and chiral phase transition with two flavor massless quarks ($N=4$) [1, 2].

The phase transition had better be investigated by finite-temperature field theory, which is based on the statistical principle only. The perturbation theory, however, breaks down around the critical temperature when the phase transition is second or weakly first order [3], even if daisy-diagrams (ring-diagrams) [4, 5, 6] are resummed. Investigation into phase transitions at finite-temperature has long been hampered by this failure. Many methods to avoid the failure were proposed: CJT method [7], renormalization improvement [8], novel summation [9], padé improvement [10], exact renormalization group with temperature [11], and auxiliary-mass method [12, 13, 14].

The auxiliary-mass method is used in the present paper. The method is based on the following idea. First an effective potential can be calculated at large mass by the perturbation theory since it is reliable there. Second the effective potential is extrapolated to the small mass range, where the perturbation theory is not reliable, using an evolution equation. Finally various quantities (e.g. critical exponents) are determined from the effective potential. The method is applied to $O(N)$ scalar model concretely in the following; the phase transition of the model is investigated and the critical exponents are determined.

We consider the following Lagrangian density,

$$\mathcal{L}_E = -\frac{1}{2} \left(\frac{\partial \phi_a}{\partial \tau} \right)^2 - \frac{1}{2} (\nabla \phi_a)^2 - \frac{1}{2} m^2 \phi_a^2 - \frac{\lambda}{4!} (\phi_a^2)^2 + J_a \phi_a + c.t. \quad , \quad (1)$$

where J_a is an external source function and the index “ a ” runs from 1 to N . First the effective potential V is calculated by the perturbation theory within one-loop order at large mass $m^2 = M^2 = O(T^2)$. We choose field expectation values $\bar{\phi}_a = \bar{\phi} \delta_{1a}$ without a loss of generality because of $O(N)$

invariance:

$$V = \frac{1}{2}M^2\bar{\phi}^2 + \frac{\lambda}{4!}\bar{\phi}^4 + \frac{T}{2\pi^2} \int_0^\infty dr r^2 \log \left[1 - \exp \left(-\frac{1}{T} \sqrt{r^2 + M^2 + \frac{\lambda}{2}\bar{\phi}^2} \right) \right] \\ + (N-1) \frac{T}{2\pi^2} \int_0^\infty dr r^2 \log \left[1 - \exp \left(-\frac{1}{T} \sqrt{r^2 + M^2 + \frac{\lambda}{6}\bar{\phi}^2} \right) \right]. \quad (2)$$

We note that the daisy-resummation is not necessary because of the large mass and one-loop zero-temperature effect is negligible if the coupling is weak. Next this effective potential is extrapolated to smaller mass using the following evolution equation [13],

$$\frac{\partial V}{\partial m^2} = \frac{1}{2}\bar{\phi}^2 + \frac{1}{4\pi i} \int_{-i\infty+\epsilon}^{+i\infty+\epsilon} dp_0 \int \frac{d^3\mathbf{p}}{(2\pi)^3} \frac{1}{-p_0^2 + \mathbf{p}^2 + m^2 + \frac{\lambda}{2}\bar{\phi}^2 + \Pi_{Sig}} \frac{1}{e^{\frac{p_0}{T}} - 1} \\ + \frac{N-1}{4\pi i} \int_{-i\infty+\epsilon}^{+i\infty+\epsilon} dp_0 \int \frac{d^3\mathbf{p}}{(2\pi)^3} \frac{1}{-p_0^2 + \mathbf{p}^2 + m^2 + \frac{\lambda}{6}\bar{\phi}^2 + \Pi_{NG}} \frac{1}{e^{\frac{p_0}{T}} - 1}. \quad (3)$$

Here $\Pi_{Sig} = \Pi_{Sig}(m^2, \bar{\phi}^2, p_0^2, \mathbf{p}^2, T)$ and $\Pi_{NG} = \Pi_{NG}(m^2, \bar{\phi}^2, p_0^2, \mathbf{p}^2, T)$ are the full self-energy, which are those for massive and massless mode in the broken phase. This equation is modified from that of $\lambda\phi^4$ theory [13] straitforwardly through a diagonalization of the propagator. Though this equation is exact, it can not be solved without an approximation; because it includes the full propagator which can not be known exactly. We then replace as follows,

$$m^2 + \frac{\lambda}{2}\bar{\phi}^2 + \Pi_{Sig}(0, 0, \bar{\phi}, m^2, \tau) \rightarrow \frac{\partial^2 V}{\partial \bar{\phi}^2} \\ m^2 + \frac{\lambda}{6}\bar{\phi}^2 + \Pi_{NG}(0, 0, \bar{\phi}, m^2, \tau) \rightarrow \frac{1}{\bar{\phi}} \frac{\partial V}{\partial \bar{\phi}}. \quad (4)$$

This replacement corresponds to fixing an external momentum of the full self-energy to zero¹. It allows us to convert eq.(3) to the following partial differential equation,

$$\frac{\partial V}{\partial m^2} = \frac{1}{2}\bar{\phi}^2 + \frac{1}{4\pi^2} \int_0^\infty dr r^2 \frac{1}{\sqrt{r^2 + \frac{\partial^2 V}{\partial \bar{\phi}^2}}} \frac{1}{\exp \left(\frac{1}{T} \sqrt{r^2 + \frac{\partial^2 V}{\partial \bar{\phi}^2}} \right) - 1}$$

¹ This is the first approximation of a systematic calculation [15].




Figure 1: The effective potential obtained by the auxiliary-mass method ($N = 4, \lambda = 1$). Second-order phase transition occurs at the critical temperature. Similar behavior are observed for the other N and λ .

$$+ \frac{N-1}{4\pi^2} \int_0^\infty dr r^2 \frac{1}{\sqrt{r^2 + \frac{1}{\phi} \frac{\partial V}{\partial \phi}}} \frac{1}{\exp\left(\frac{1}{T} \sqrt{r^2 + \frac{1}{\phi} \frac{\partial V}{\partial \phi}}\right) - 1}. \quad (5)$$

We numerically solve the equation to $m^2 = -\mu^2$, where $O(N)$ symmetry is broken at zero-temperature, under the initial condition eq.(2). We solve it above the critical temperature only and can get sufficient information about the critical phenomenon.

The effective potential are calculated using numerical methods in Ref.[14]. They are shown in fig.1. We can not observe a second minimum which is a hallmark of a first order phase transition. Instead, the curvature of the origin decreases and vanishes smoothly as temperature decreases; one can, therefore, observe that the phase transition of the model is second order as it should be.

The critical temperature is shown as a function of N in fig.2. Though they resemble to a result of the perturbation theory at leading order [16], $T_c = 6\sqrt{\frac{2}{\lambda(N+2)}}$, which is determined from the condition that mass with the daisy-diagram vanishes, they have a slight difference quantitatively.




Figure 2: Critical temperature as a function of N at $\lambda = 1$ ($\diamond$). This resembles to a result of the perturbation theory at leading order (—) but has a slight difference quantitatively.

Finally the critical exponents, γ and δ , are determined using this potential. We determine γ from a second derivative of the effective potential with respect to $\bar{\phi}$ at the origin. The critical exponent γ is defined as follows,

$$\chi \equiv \left. \frac{\partial \bar{\phi}}{\partial J_1} \right|_{J_a=0} \sim \chi \tau^{-\gamma} \quad (\tau = (T - T_c)/T_c) \quad (6)$$

the following identity then relates the second derivative to the susceptibility,

$$\left. \frac{\partial \bar{\phi}}{\partial J_1} \right|_{J_a=0} = \left(\frac{\partial^2 V}{\partial \bar{\phi}^2} \right)^{-1} \Big|_{\bar{\phi}=\phi_c}. \quad (7)$$

Here ϕ_c is determined from the condition, $\frac{\partial V}{\partial \bar{\phi}} \Big|_{\bar{\phi}=\phi_c} = 0$. The second derivative are shown in log-scale in fig.3. We then determine the gradients, which is the very γ we want. One can observe that they become steeper as N increases; then, γ become larger. The results are $\gamma = 1.13$ ($N \rightarrow 0$), 1.37 ($N = 1$), 1.47 ($N = 2$), 1.60 ($N = 3$), 1.66 ($N = 4$). These results are summarised in table.(1), compared with the Landau approximation and world best values.




Figure 3: Second derivative of the effective potential with respect to ϕ . The gradients are steeper for larger N .

We determine δ using the effective potential at the critical temperature. The critical exponent δ is defined as follows.

$$\bar{\phi} \propto J_1^{1/\delta} = \left(\frac{\partial V}{\partial \phi}\right)^{1/\delta} \quad (T = T_c). \quad (8)$$

The following relation, which derived from eq.(8), enables us to determine δ from the effective potential at the critical temperature,

$$V \propto \bar{\phi}^{\delta+1} \quad (T = T_c). \quad (9)$$

The effective potential at the critical temperature is shown in fig.1. We determine δ from it. The result are $\delta = 3.8$ ($N \rightarrow 0$), 4.0 ($N = 1$), 4.2 ($N = 2$), 4.4 ($N = 3$), 4.4 ($N = 4$).

In the present paper, we showed that the phase transition of $O(N)$ scalar model is second order using auxiliary-mass method at finite temperature. This is a great progress because we can not show it using the perturbation theory with daisy-resummation — the traditional method to calculate the effective potential at finite temperature — [3].

Since the phase transition turned out to be second order, we determined the critical exponents of the phase transition. Though the results are not as

	γ (LA,WBV)	δ (LA,WBV)
N $\rightarrow$ 0 [1]	1.13 (1, 1.16)	3.8 (3, 4.77)
N=1 [1]	1.37 (1, 1.24)	4.0 (3, 4.76)
N=2 [1]	1.47 (1, 1.32)	4.2 (3, 4.78)
N=3 [1]	1.60 (1, 1.40)	4.4 (3, 4.78)
N=4 [18]	1.66 (1, 1.48)	4.4 (3, 4.85)

Table 1: The critical exponents, γ and δ , obtained in the present paper. Those of Landau approximation (LA) and world best values (WBV) are also summarised. We used lattice results as WBV.

accurate as world best values, they are much better than that obtained in the Landau approximation. The error would be due to the replacement (4). We must improve the approximation in order to get more accurate values [15].

In conclusion, the auxiliary-mass method turns out to be reliable not only qualitatively but also in rough quantitative estimation in $O(N)$ invariant scalar model. This method would be reliable in the other models. What is more, we can investigate not only second-order phase transition but also first-order phase transition without any modifications [17]. This method, therefore, enable us to investigate the phase transition of various models: the cubic anisotropy model, the abelian Higgs model, and the standard model.

We finally express our thanks to T. Inagaki for valuable discussions and communications. J.S. is supported by JSPS fellowship.

References

- [1] J. Zinn-Justin, *Quantum Field Theory and Critical Phenomena*, (Oxford Univ. Press, 1996).
- [2] R.Pisarski and F.Wilczek, *Phys.Rev.* **D29** (1984) 338.
- [3] P. Arnold and O. Espinosa, *Phys. Rev.* **D47** (1993) 3546.

- [4] J. I. Kapsta, *Finite Temperature Field Theory* (Cambridge Univ. Press, 1989).
- [5] K.Takahashi, *Z.Phys.***C26** (1985) 601.
- [6] P. Fendley, *Phys. Lett.* **B196** (1987) 175.
- [7] G. Amelino-Camelia, *Phys.Lett.* **B407** (1997) 268.
- [8] H.Nakkagawa and H.Yokota, *Mod.Phys.Lett.* **A11** (1996) 2259.
- [9] S. Chiku and T. Hatsuda, *Phys.Rev.* **D57** (1998) 6.
- [10] T. Hatsuda, *Phys.Rev.* **D56** (1997) 8111.
- [11] S.B. Liao and M.Strickland, *Phys.Rev.* **D52** (1995) 3653,*Nucl.Phys. B497* **B497** (1997) 611.
- [12] I. T. Drummond, R. R. Horgan, P. V. Landshoff and A. Rebhan, *Phys. Lett.* **B398** (1997) 326.
- [13] T.Inagaki, K.Ogure and J.Sato, hep-th/9705133.
- [14] K.Ogure and J.Sato, hep-ph/9801439.
- [15] K.Ogure and J.Sato, in preparation.
- [16] L. Dolan and R. Jackiw, *Phys.Rev.* **D9** (1974) 3320.
- [17] J.Arafune, K.Ogure and J.Sato, *Prog.theor.Phys.* **99** (1998) 119.
- [18] K.Kanaya and S.Kaya *Phys.Rev.* **D51** (1995) 2404.